
Using Shuttle Radar Topography Mission Elevation Data to Map Mangrove Forest Height in the Caribbean




Marc Simard

(email: <u>marc.simard@jpl.nasa.go</u>v)

Caltech - NASA - Jet Propulsion Laboratory K. Zhang(FIU), V. H. Rivera-Monroy(LSU), M. Ross (FIU)




Mangrove Distribution





- Biodiversity
 - Habitats of 1300 species of animals
 - 628 mammals, birds, reptiles, fish and amphibians
- o Among the most productive ecosystems on earth
 - o 170k km² with mean 2.5g C m-² per day (10t/ha/yr)



- Mangrove Sediment
- Recycled
- Export of Ocean














- Biodiversity
 - o Habitats of 1300 species of animals
 - o 628 mammals, birds, reptiles, fish and amphibians
- Among the most productive ecosystems on earth
 - $_{\circ}$ 170k km² with mean 2.5g C m⁻² per day (10t/ha/yr)



- Mangrove Sediment
- Recycled
- Export of Ocean
- o Annual input into ocean 46X1012 g C
 - o Thus contributes 11% of global total export to ocean










- Biodiversity
 - o Habitats of 1300 species of animals
 - o 628 mammals, birds, reptiles, fish and amphibians
- Among the most productive ecosystems on earth
 - \circ 170k km² with mean 2.5g C m⁻² per day (10t/ha/yr)



- Mangrove Sediment
 - Recycled
- Export of Ocean
- Annual input into ocean 46X1012 g C
 - o Thus contributes 11% of global total export to ocean



Mangroves


Forests





- Biodiversity
 - o Habitats of 1300 species of animals
 - o 628 mammals, birds, reptiles, fish and amphibians
- Among the most productive ecosystems on earth
 - \circ 170k km² with mean 2.5g C m⁻² per day (10t/ha/yr)



- Mangrove Sediment
- Recycled
- Export of Ocean
- o Annual input into ocean 46X1012 g C
 - o Thus contributes 11% of global total export to ocean











- **Biodiversity**
 - Habitats of 1300 species of animals
 - 628 mammals, birds, reptiles, fish and amphibians
- Among the most productive ecosystems on earth
 - 170k km² with mean 2.5g C m-2 per day (10t/ha/yr)



- Mangrove Sediment
- Recycled
- **Export ot Ocean**
- Annual input into ocean 46X1012 g C
 - Thus contributes 11% of global total export to ocean









- Biodiversity
 - Habitats of 1300 species of animals
 - 628 mammals, birds, reptiles, fish and amphibians
- Among the most productive ecosystems on earth
 - o $170k \text{ km}^2 \text{ with mean } 2.5g \text{ C m}^{-2} \text{ per day } (10t/\text{ha/yr})$





Mangrove Sediment


Recycled

Export ot Ocean

- Annual input into ocean 46X1012 g C
 - Thus contributes 11% of global total export to ocean










- Biodiversity
 - o Habitats of 1300 species of animals
 - o 628 mammals, birds, reptiles, fish and amphibians
- Among the most productive ecosystems on earth
 - \circ 170k km² with mean 2.5g C m⁻² per day (10t/ha/yr)



- Mangrove Sediment
- Recycled
- Export of Ocean
- o Annual input into ocean 46X1012 g C
 - Thus contributes 11% of global total export to ocean








- Biodiversity
 - Habitats of 1300 species of animals
 - o 628 mammals, birds, reptiles, fish and amphibians
- Among the most productive ecosystems on earth
 - o 170k km² with mean 2.5g C m-² per day (10t/ha/yr)




- Mangrove Sediment
- Recycled
- Export of Ocean
- o Annual input into ocean 46X1012 g C
 - o Contributing 11% of global total export to ocean




• Contributing 15% of carbon accumulation in modern sediments.








Mangroves



- Biodiversity
 - Habitats of 1300 species of animals
 - o 628 mammals, birds, reptiles, fish and amphibians
- Among the most productive ecosystems on earth
 - $_{\circ}$ 170k km² with mean 2.5g C m⁻² per day (10t/ha/yr)



- Mangrove Sediment
- Recycled
- Export of Ocean
- o Annual input into ocean 46X1012 g C
 - o Contributing 11% of global total export to ocean



• Contributing 15% of carbon accumulation in modern sediments.




87%

Forests




Mangroves

- Biodiversity
 - Habitats of 1300 species of animals
 - o 628 mammals, birds, reptiles, fish and amphibians
- Among the most productive ecosystems on earth
 - $_{\circ}$ 170k km² with mean 2.5g C m⁻² per day (10t/ha/yr)



- Mangrove Sediment
- Recycled
- Export of Ocean
- o Annual input into ocean 46X1012 g C
 - o Contributing 11% of global total export to ocean




• Contributing 15% of carbon accumulation in modern sediments.










Mangroves Are Endangered

- Endangered by Urbanization, exploitation and sea level rise
- Already 35% of mangrove forests have disappeared and 60% could be lost by 2030 (UNEP report 2006);
- The estimated economical value varies between \$200k to \$900k per km² per year (UNEP report 2006);



Shuttle Radar Topography Mission (SRTM)




- 1. Radar Interferometer
- 2. 60m boom
- 3. C and X band
- 4. 11 days in February 2000.
- 5. 80% Earth Land
- 6. Latitude 56S to 60N
- 7. 30m US and World 90m

http://www2.jpl.nasa.gov/srtm/




SRTM Elevation data








Relative Vegetation Height





LIDAR Data





- o Optech 1233-infrared 1um
- First and last reflection method (No Waveform)
- o Nominal Altitude of 500m
 - o 360m swath
- o May 13-15 2004
- 1.5m spacing with 13cm laser footprint
- o 15 cm elevation accuracy






Lidar Tansects









Results Everglades National Park

Mapping Mangrove Forest Height in the Everglades National Park (ENP) Using SRTM




Marc Simard et al., "Mapping Height and Biomass of Mangrove Forests in Everglades National Park with SRTM Elevation Data", *Photogrammetric Engineering and Remote Sensing*, SRTM special issue, April 2006

Mapping Biomass of Mangrove Forest in the Everglades National Park (ENP) using SRTM



Global storage of Carbon in Mangrove Biomass* is (Twilley et al 1992): 4.03 x 10¹⁵ grams Current first-rate estimation of carbon storage in Mangroves in the ENP is: 2.52 x 10¹² grams (45%C DM)

How to apply this methodology worldwide?




- Extrapolate results from other regions
- Use ICEsat data
- Collect field data


ICEsat/GLAS

Geoscience Laser Altimeter System

- 172m spacing
- 64m footprint



ICEsat/GLAS Waveform



California Institute of Technology

ICEsat Waveform:

Characterizing Structure


Degraded mangrove waveform

- Multiple Gaussians to fit ground, high and low canopies
- Compute waveform centroid


Tall mangrove waveform

Compute canopy waveform centroid







Field Data




Forest height estimation

Cienaga Grande de Santa Marta, Colombia Calibrated with ICEsat LIDAR Waveforms






Cienaga Grande de Santa Marta, Colombia Calibrated with ICEsat LIDAR Waveforms




Landsat Land Cover Classification




Mean Mangrove Tree Height

San Juan River Venezuela


SRTM C- and X-band






Mangrove height (0-25m) Difference X-C from 0.6 TO 3.5m





Shrimp Farming Development in Honduras

Shrimp Farming in the Southern Region of the Gulf of Fonseca



Estimated total area of shrimp ponds:

1985 = 845 ha


1999 = 15,580 ha

(V. H. Rivera-Monroy, U. Louisiana at Lafayette)



Granjas Marinas San Bernardo Shrimp Farm (7,000 ha); Largest shrimp farm the Gulf of Fonseca.

AIRSAR Vegetation Height Estimate (Chismuyo, Gulf of Fonseca)





Conclusions

- SRTM provides a systematic tool to map mangrove height and biomass
- o The method is simple and works well for mangrove forests.
- ICEsat shows great potential to calibrate SRTM data and identify a variety of forest structures
- Need to apply the method to other mangrove forests around the globes
- Need to derive a relation between ICEsat waveform and canopy structure parameters
- o Quantify the Impact of Hurricanes on Mangrove Forest



Hurricane impact

Two Category 1 hurricanes Katrina august 25th, 2005 Wilma October 24th, 2005




Hurricane Damage











<VALUE>

-4.963091373 - -2.782808766

___-2.782808765 - -1.420132136

-1.420132135 - -0.602526158

-0.602526158 - -0.329990832

-0.329990832 - 0.215079820

0.215079820 - 1.305221124

1,305221125 - 2,395362428

2.395362429 - 3.485503732

3,485503733 - 4,575645035

4,575645036 - 5,665786339

5.66578634 - 6.755927643

6.755927644 - 7.846068947

7.846068948 - 8.936210251

8.936210252 - 10.02635155

10.02635156 - 11.11649286

<u>11.11649287 - 11.93409884</u>

11.93409885 - 13.02424014

13.02424015 - 14.11438144

14.11438145 - 15.20452275

15,20452276 - 16,02212873

16.02212874 - 16.8397347

16.83973471 - 17.65734068

<u>17.65734069 - 18.47494666</u>

18.47494667 - 19.29255264

19.29255265 - 20.11015862


20.11015863 - 20.92776459

20.9277646 - 21.74537057


21.74537058 - 22.83551188

22.83551189 - 26.10593579

26.1059358 - 64.53341675



Tree height histograms 2004 and 2006





Elevation Histograms for First Returns of Tile 1 (T1)

2006

X axis: elevation (m, NAVD88)

Y axis: number of Lidar points





Conclusions

- What is the mean tree height?
- Can we understand structure from LIDAR data?
- How to quantify the impact of hurricanes on mangrove forests using active remote sensing?